



# Variational Iteration Method for Fredholm integral equations of the second kind

**J. Biazar<sup>a,\*</sup>, H. Ebrahimi<sup>b</sup>**

<sup>a</sup> *Department of Mathematics, Faculty of Science, University of Guilan,*

<sup>b</sup> *Department of Mathematics, Islamic Azad University, Rasht branch.*

---

## Abstract

In this paper, He's variational iteration method is applied to Fredholm integral equations of the second kind. To illustrate the ability and simplicity of the method, some examples are provided. The results reveal that the proposed method is very effective and simple and for first four examples leads to the exact solution.

**Keywords:** Variational iteration method; Fredholm integral equation; Lagrange multiplier; Restricted variation.

---

---

\*Corresponding author, E-mail: [jafai.biazar@gmail.com](mailto:jafai.biazar@gmail.com)(J. Biazar)

## 1 Introduction

Variational iteration method [1-2] is a power device for solving various kinds of linear and non-linear functional equations. It was introduced by Ji-Huan He in 1998 and has been used by many mathematicians and engineers to solve various kinds of functional equations, such as wave equation [9], hyperbolic differential equations [10], Telegraph equation [11], nonlinear chemistry problems [12], Cauchy reaction diffusion problem [13], quadratic Riccati differential equation [14], and many other equations. Fredholm integral equation has been solved by some other methods, such as Adomian decomposition method [3, 7] and Homotopy perturbation method [3-6]. In this study, we use variational iteration method for fredholm integral equations of the second kind. The general form of this integral equation is given by

$$u(x) = f(x) + \int_a^b k(x,t) F(u(t)) dt, \quad a \leq x \leq b. \quad (1)$$

Where  $k(x,t)$  is the kernel of the integral equation,  $f$  and  $F$  are known functions and  $u(x)$  is the unknown solution of integral equation, which we are going to find, via variational iteration method.

## 2 Application of Variational iteration method

For solving equation (1) by variational iteration method, first we differentiate once from both sides of equation (1) with respect to  $x$  :

$$u'(x) = f'(x) + \int_a^b \frac{\partial k(x,t)}{\partial x} F(u(t)) dt. \quad (2)$$

Now, we apply variational iteration method for equation (2). According to this method correction functional can be written in the following form:

$$u_{n+1}(x) = u_n(x) + \int_0^x \lambda(s) \left( u'_n(s) - f'(s) - \int_a^b \frac{\partial k(s,t)}{\partial s} F(\tilde{u}_n(t)) dt \right) ds, \quad (3)$$

where  $\lambda(s)$  is a general Lagrange multiplier. To make the above correction functional stationary with respect to  $u_n$ , we have:

$$\begin{aligned} \delta u_{n+1}(x) &= \delta u_n(x) + \delta \int_0^x \lambda(s) \left( u'_n(s) - f'(s) - \int_a^b \frac{\partial k(s,t)}{\partial s} F(\tilde{u}_n(t)) dt \right) ds \\ &= \delta u_n(x) + \int_0^x \lambda(s) \delta \left( u'_n(s) \right) ds = \delta u_n(x) + \lambda(x) \delta u_n(x) + \int_0^x \lambda'(s) \delta u_n(s) ds = 0. \end{aligned}$$

From the above relation for any  $\delta u_n$ , we obtain the Euler-Lagrange equation

$$\lambda'(s) = 0. \quad (4)$$

With the following natural boundary condition:

$$\lambda(x) + 1 = 0. \quad (5)$$

By using equations (4) and (5), Lagrange multiplier can be identified optimally as follows:

$$\lambda(s) = -1. \quad (6)$$

Substituting the identified Lagrange multiplier into equation (3), results in the following iterative formula:

$$u_{n+1}(x) = u_n(x) + \int_0^x \lambda \left( u'_n(s) - f'(s) - \int_a^b \frac{\partial k(s,t)}{\partial s} F(u_n(t)) dt \right) ds. \quad (7)$$

By starting from  $u_0(x)$ , we can obtain the exact solution or an approximate solution of the equation (1). In some Fredholm integral equations by differentiating from the integral equation, we obtain a differential equation and we can solve this differential equation by variational iterative method. (See Example 2)

### 3 Numerical examples

To illustrate the ability and simplicity of the proposed technique, some examples are provided here.

**Example 1.** Consider the following linear Fredholm integral equation

$$u(x) = e^{3x} - \frac{1}{9}(2e^3 + 1)x + \int_0^1 x t u(t) dt, \quad 0 \leq x \leq 1, \quad (8)$$

with the exact solution,  $u(x) = e^{3x}$  [3, 4].

The corresponding iterative formula (7) for this example can be constructed as follows:

$$u_{n+1}(x) = u_n(x) - \int_0^x \left( u'_n(s) - 3e^{3s} - \frac{1}{9}(2e^3 + 1) - \int_0^1 t u_n(t) dt \right) ds. \quad (9)$$

By taking  $u_0(x) = e^{3x} - \frac{1}{9}(2e^3 + 1)x$ , we derive the following results:

$$u_1(x) = e^{3x} - \frac{x}{3^3}(2e^3 + 1),$$

$$u_2(x) = e^{3x} - \frac{x}{3^4}(2e^3 + 1),$$

$$u_3(x) = e^{3x} - \frac{x}{3^5} (2e^3 + 1),$$

⋮

$$u_n(x) = e^{3x} - \frac{x}{3^{n+2}} (2e^3 + 1),$$

⋮

Thus, we have  $u(x) = \lim_{n \rightarrow \infty} u_n(x) = e^{3x}$ , which is the exact solution.

**Example 2.** Consider the following fredholm integral equation with the exact solution  $u(x) = \sin x + \cos x$  [4].

$$u(x) = (1 - 2\pi) \cos x + \sin x + 4 \int_0^\pi \cos x \cos t u(t) dt. \quad (10)$$

By differentiating twice from integral equation (10), the following differential equation will be obtained:

$$u''(x) + u(x) = 0. \quad (11)$$

By applying variational iterative method for (11), we derive the following iterative formula:

$$u_{n+1}(x) = u_n(x) + \int_0^x \sin(s-x) (u_n''(s) + u_n(s)) dt. \quad (12)$$

Let's start with  $u_0(x) = A \sin x + B \cos x$ , as an initial approximation, and using the iterative formula (12) we get:

$$u_1(x) = A \sin x + B \cos x,$$

$$u_2(x) = A \sin x + B \cos x,$$

$$\vdots$$

For determining  $A$  and  $B$ , we substitute the solution in the integral equation (10) and we obtain  $A = B = 1$ .

**Example 3.** Consider the following integral equation with the exact solution,  $u(x) = x$  [8].

$$u(x) = \frac{7}{8}x + \frac{1}{2} \int_0^1 x t u^2(t) dt. \tag{13}$$

The iterative formula would be as:

$$u_{n+1}(x) = u_n(x) - \int_0^x \left( u'_n(s) - \frac{7}{8} + \frac{1}{2} \int_0^1 t u_n^2(t) dt \right) ds. \tag{14}$$

Consider the initial approximation  $u_0(x) = \frac{7}{8}x$ . Therefore other terms of the sequence are computed as follows:

$$u_1(x) = \frac{497}{512}x = 0.9707031250x,$$

$$u_2(x) = \frac{2082017}{2097152}x = 0.9927830696x,$$

$$u_3(x) = 0.9982022779x,$$

$$u_4(x) = 0.9995509734x,$$

$$u_5(x) = 0.9998877686x,$$

$\vdots$

The sequence tends to  $x$ , as  $n \rightarrow +\infty$ .

**Example 4.** Consider the following equation

$$u(x) = \cos x - x + \int_0^1 x(u^2(t) - \sin^2 t) dt, \tag{15}$$

with the exact solution,  $u(x) = \cos x$  [6].

In the same procedure, the iterative formula can be expressed as the following:

$$u_{n+1}(x) = u_n(x) - \int_0^x \left( u'_n(s) + \sin s + 1 - \int_0^1 (u_n^2(t) - \sin^2 t) dt \right) ds. \tag{16}$$

By using this iterative formula and taking  $u_0(x) = \cos x$ , we have:

$$u_1(x) = \cos x,$$

$$u_2(x) = \cos x,$$

⋮

Therefore, the exact solution can be recognized easily.

**Example 5.** For the last example consider the following non-linear Fredholm integral equation

$$u(x) = -\sin(4x) - x^3 \left( -\frac{367}{4096} \cos(4) \sin(4) + \frac{11357}{98304} - \frac{2095}{32768} \cos^2(4) \right) + \int_0^1 x^3 t^5 u^2(t) dt. \tag{17}$$

Where the exact solution is,  $u(t) = \sin(-4x)$  [5].

By using variational iteration method for equation (17) and considering


$$u_0(x) = -\sin(4x) - x^3 \left( -\frac{367}{4096} \cos(4) \sin(4) + \frac{11357}{98304} - \frac{2095}{32768} \cos^2(4) \right),$$

we obtain the following results:

$$\begin{aligned} u_1 &= -0.003982967666 x^3 - 8. \sin(x) \cos(x)^3 + 4. \sin(x) \cos(x) , \\ u_2 &= -0.000374692464 x^3 - 1. \sin(4. x) , \\ u_3 &= -0.0000353613617 x^3 - 1. \sin(4. x) , \\ u_4 &= -0.333819270 \cdot 10^{-5} x^3 - 1. \sin(4. x) , \\ u_5 &= -0.315123700 \cdot 10^{-6} x^3 - 1. \sin(4. x) , \\ &\vdots \end{aligned}$$

Suppose  $u(x) \approx u_5(x)$ . Plots of approximated solution,  $u_5(x)$ , and the exact solution are presented in Fig.1.

Fig. 1: Plots of the exact and approximate solutions for Example 5



#### 4 Conclusion and Discussion



In this work, variational iterative method has been successfully applied to find the solution of Fredholm integral equations of the second kind. It can be concluded that the method is very powerful and efficient technique for finding exact solutions for wide classes of problems. In this work, we have used the Maple 11 package to carry out the computations.

## **References**

- [1] He, J. H., Variational iteration method a kind of non-linear analytical technique: some examples, *International Journal of Non-Linear Mechanics*, 34(4), 699-708, 1999.
- [2] He, J. H., Variational iteration method-Some recent results and new interpretations, *Journal of Computational and Applied Mathematics*, 207, 3-17, 2007.
- [3] Abbasbandy, S., Numerical solutions of the integral equations: Homotopy perturbation method and Adomian's decomposition method, *Applied Mathematics and Computation*, 173, 493-500, 2006.
- [4] Golbabai, A., and Keramati, B., Modified homotopy perturbation method for solving Fredholm integral equations, *Chaos, Solitons and Fractals*, 2006 in press.
- [5] Javidi, M., and Golbabai, A., Modified homotopy perturbation method for solving non-linear Fredholm integral equations, *Chaos, Solitons and Fractals*, 2007 in press.
- [6] Ganji, D. D., Afrouzi, G. A., Hosseinzadeh, H., and Talarposhti, R. A., Application of homotopy-perturbation method to the second kind of the nonlinear integral equations, *Physics Letters A*, 371, 20-25, 2007.

- [7] Taghizadeh, N., and Ebrahimi, H., The solution of Fredholm integral equations with degenerate kernel by Adomian decomposition method, *Korean Annals of Math*, 22, NO. 2, 163-172, 2005.
- [8] Jerri, A. J., *Introduction to integral equation with applications*, 2th ed., John Wily and Sons, Inc., New York, 1999.
- [9] Biazar, J., and Ghazvini, H., An analytical approximation to the solution of a wave equation by a Variational iteration method, *Applied Mathematics Letters*, 21 (8), 780-785, 2008.
- [10] Biazar, J., and Ghazvini, H., He's Variational iteration method for solving hyperbolic differential equations, *International Journal of nonlinear sciences and numerical simulation* 8 (3), 311-314, 2007.
- [11] Biazar, J., and Ebrahimi, H., An Approximation to the Solution of Telegraph Equation by Variational Iteration Method, *Numerical Methods for Partial Differential Equations*, In press.
- [12] Ganji, D. D., Nourollahi, M., and Mohseni, E., Application of He's methods to nonlinear chemistry problems, *Computers and Mathematics with applications*, 54 (7-8), 1122-1132, 2007.
- [13] Dehghan, M., and Shakeri, F., Application of He's Variation iteration method for solving the Cauchy reaction diffusion problem, *Journal of computational and Applied Mathematics*, 217 (2), 435-446, 2008.
- [14] Abbasbandy, S., A new application of He's Variational iteration method for quadratic Riccati differential equation by using Adomian's polynomials, *Journal of Computational and Applied Mathematics*, 207(1), 59-63, 2007.