



## Solution of fuzzy differential equations under generalized differentiability by Adomian decomposition method

**T. Allahviranloo\*, L. Jamshidi**

*Department of Mathematics, Science and Research Branch, Islamic Azad  
University, Tehran, 14778, Iran*

---

### Abstract

Adomian decomposition method has been applied to solve many functional equations so far. In this article, we have used this method to solve the fuzzy differential equation under generalized differentiability. We interpret a fuzzy differential equation by using the strongly generalized differentiability. Also one concrete application for ordinary fuzzy differential equation with fuzzy input data are given.

**Keywords:** Adomian decomposition method; generalizations of the differentiability of fuzzy differential equations

---

---

\*Corresponding author, E-mail: [tofigh@allahviranloo.com](mailto:tofigh@allahviranloo.com)

## 1 Introduction

Usage of fuzzy differential equations is a natural way to model dynamical systems under possibilistic uncertainty. Tainty [1]. Strongly generalized differentiability was introduced in [2] and studied in [3]. we use this differentiability concept in the present paper. Adomian decomposition method has a useful feature that provides the solution in a rapid convergent power series with elegantly computable convergence of the solution.

This paper is organized as follows: In section 2 we bring some basic definitions of fuzzy subsets and distance between fuzzy numbers. In section 3 we define a fuzzy differential equations under generalized differentiability and In section 4 we discuss Adomian decomposition method. In section 5, we have one example of this method and the conclusion and future research is drawn in section 6.

## 2 Preliminaries

We begin this section with defining the notation we will use in the paper. Let us denote by  $\mathfrak{R}_f$  the class of fuzzy subsets of the real axis  $u : \mathfrak{R} \rightarrow [0,1]$ , satisfying the following properties:

- (i)  $u$  is normal, i.e.  $\exists x_0 \in \mathfrak{R}$  with  $u(x_0) = 1$ ;
- (ii)  $u$  is convex fuzzy set  
(i.e.  $u(tx + (1-t)y) \geq \min\{u(x), u(y)\}, \forall t \in [0,1], x, y \in \mathfrak{R}$ );
- (iii)  $u$  is upper semi continuous on  $\mathfrak{R}$ ;
- (iv)  $\{x \in \mathfrak{R} : u(x) > 0\}$  is compact, where  $\bar{A}$  denotes the closure of  $A$ .

Then  $\mathfrak{R}_f$  is called the space of fuzzy numbers (see e.g. [4]). Obviously  $\mathfrak{R} \subset \mathfrak{R}_f$ .

Here  $\mathfrak{R} \subset \mathfrak{R}_f$  is understood as  $\mathfrak{R} = \{\chi_{\{x\}}; x \text{ is usual real number}\}$ . For  $0 < r \leq 1$ , denote  $[u]^r = \{x \in \mathfrak{R}; u(x) \geq r\}$  and  $[u]^0 = \{x \in \mathfrak{R}; u(x) > 0\}$ . Then

it is well-known that for any  $r \in [0,1], [u]^r$  is a bounded closed interval.

For  $u, v \in \mathfrak{R}_f$ , and  $\lambda \in \mathfrak{R}$ , the sum  $u + v$  and the product  $\lambda.u$  are defined by

$$[u + v]^r = [u]^r + [v]^r,$$

$$[\lambda.u]^r = \lambda[u]^r, \forall r \in [0,1],$$

Where

$[u]^r + [v]^r = \{x + y : x \in [u]^r, y \in [v]^r\}$  means the usual addition of two intervals (subsets) of  $\mathfrak{R}$  and  $\lambda[u]^r = \{\lambda x : x \in [u]^r\}$  means the usual product between a scalar and a subset of  $\mathfrak{R}$  (see e.g. [4,5]).

Let  $D : \mathfrak{R}_f \times \mathfrak{R}_f \rightarrow \mathfrak{R}_+ \cup \{0\}, D(u, v) = \sup_{r \in [0,1]} \max\{u_-^r - v_-^r, u_+^r - v_+^r\}$  be the Hausdorff distance between fuzzy numbers, where  $[u]^r = [u_-^r, u_+^r], [v]^r = [v_-^r, v_+^r]$ . The following properties are well-known (see e.g. [5,6])

$$D(u + w, v + w) = D(u, v), \forall u, v, w \in \mathfrak{R}_f,$$

$$D(k.u, k.v) = |k|D(u, v), \forall k \in \mathfrak{R}, u, v \in \mathfrak{R}_f,$$

$$D(u + v, w + e) \leq D(u, w) + D(v, e), \forall u, v, w, e \in \mathfrak{R}_f$$

and  $(\mathfrak{R}_f, D)$  is a complete metric space.

Also are known the following results and concepts.

**Definition 1.** Let  $F : (a, b) \rightarrow \mathfrak{R}_f$  and  $t_0 \in (a, b)$ . We say that F is differentiable at  $t_0$ ; if we have 2 forms as follows:

- 1- it exists an element  $F'(t_0) \in \mathfrak{R}_f$  such that, for all  $h > 0$  sufficiently near to 0, there are  $F(t_0 + h) - F(t_0), F(t_0) - F(t_0 - h)$  and the limits:

$$(1) \quad \lim_{h \rightarrow 0^+} \frac{F(t_0 + h) - F(t_0)}{h} = \lim_{h \rightarrow 0^+} \frac{F(t_0) - F(t_0 - h)}{h} = F'(t_0)$$

2- It exists an element  $F'(t_0) \in \mathfrak{R}_f$  such that, for all  $h < 0$  sufficiently near to 0, there are

$$(2) \quad \lim_{h \rightarrow 0^-} \frac{F(t_0 + h) - F(t_0)}{h} = \lim_{h \rightarrow 0^-} \frac{F(t_0) - F(t_0 - h)}{h} = F'(t_0)$$

**Theorem 2.1** Let  $F : (a, b) \rightarrow \mathfrak{R}_f$  be a function and denote  $[F(t)]^\alpha = [f_\alpha(t), g_\alpha(t)]$ , for each  $\alpha \in [0, 1]$ . Then

(i) If  $F$  is differentiable in the first form (1), then  $f_\alpha$  and  $g_\alpha$  are differentiable functions and

$$(3) \quad [F'(t)]^\alpha = [f'_\alpha(t), g'_\alpha(t)]$$

(ii) If  $F$  is differentiable in the second form (2), then  $f_\alpha$  and  $g_\alpha$  are differentiable functions and

$$(4) \quad [F'(t)]^\alpha = [g'_\alpha(t), f'_\alpha(t)]$$

Proof: See [7].

**Definition 2.** (see e.g. [8]). Let  $x, y \in \mathfrak{R}_f$ . If there exists  $z \in \mathfrak{R}_f$  such that  $x = y \oplus z$ , then  $z$  is called the H-difference of  $x$  and  $y$  and it is denoted by  $x - y$ .

**Definition 3.** (See e.g. [8].) A function  $f : (a, b) \rightarrow \mathfrak{R}_f$  is called H-differentiable on  $x_0 \in (a, b)$  if for

$h > 0$  sufficiently small there exist the H-differences  $f(x_0 + h) - f(x_0)$ ,  $f(x_0) - f(x_0 - h)$  and an element  $f'(x_0) \in \mathfrak{R}_f$  such that

$$(5) \quad 0 = \lim_{h \rightarrow 0} D\left(\frac{f(x_0 + h) - f(x_0)}{h}, f'(x_0)\right) = \lim_{h \rightarrow 0} D\left(\frac{f(x_0) - f(x_0 - h)}{h}, f'(x_0)\right)$$

(Here  $h$  at denominator means  $\frac{1}{h} \otimes .$ )

**Definition 4.** Let  $f : (a, b) \rightarrow \mathfrak{R}_f$  and  $x_0 \in (a, b)$ . We say that  $f$  is strongly generalized differentiable at  $x_0$ , if there exists an element  $f'(x_0) \in \mathfrak{R}_f$ , such that

(i) for all  $h > 0$  sufficiently small,  $\exists f(x_0 + h) - f(x_0)$ ,  $f(x_0) - f(x_0 - h)$  and the limits (in the metric  $D$ )

$$(6) \quad \lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h} = \lim_{h \rightarrow 0} \frac{f(x_0) - f(x_0 - h)}{h} = f'(x_0)$$

or

(ii) for all  $h > 0$  sufficiently small,  $\exists f(x_0) - f(x_0 + h)$ ,  $f(x_0 - h) - f(x_0)$  and the limits

$$(7) \quad \lim_{h \rightarrow 0} \frac{f(x_0) - f(x_0 + h)}{(-h)} = \lim_{h \rightarrow 0} \frac{f(x_0 - h) - f(x_0)}{(-h)} = f'(x_0)$$

or

(iii) for all  $h > 0$  sufficiently small,  $\exists f(x_0 + h) - f(x_0)$ ,  $f(x_0) - f(x_0 - h)$  and the limits

$$(8) \quad \lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h} = \lim_{h \rightarrow 0} \frac{f(x_0 - h) - f(x_0)}{(-h)} = f'(x_0)$$

or

(iv) for all  $h > 0$  sufficiently small,  $\exists f(x_0) - f(x_0 + h), f(x_0) - f(x_0 - h)$  and the limits

$$(9) \quad \lim_{h \rightarrow 0} \frac{f(x_0) - f(x_0 + h)}{(-h)} = \lim_{h \rightarrow 0} \frac{f(x_0) - f(x_0 - h)}{h} = f'(x_0)$$

### 3 A fuzzy differential equation under generalized differentiability

In this section we will study solutions to

$$(1) \quad y^{(n)} + a_{n-1}(x)y^{(n-1)} + \dots + a_1(x)y^{(1)} + a_0(x)y = g(x)$$

Where the  $a_i(x), 0 \leq i \leq n-1$ , and  $g(x)$  are continuous on some interval  $I$ , subject to initial conditions  $y(0) = \bar{\gamma}_0, y^{(1)}(0) = \bar{\gamma}_1, \dots, y^{(n-1)}(0) = \bar{\gamma}_{n-1}$ , for fuzzy numbers  $\bar{\gamma}_i, 0 \leq i \leq n-1$ , the interval  $I$  can be  $[0, T]$  for some  $T > 0$  or  $I = [0, \infty)$  by using the derivative in strongly generalized sense.

Consider three case while coefficient are positive and/or negative.

**Case 1:**  $a_i \geq 0, i = 0, \dots, n-1$

(i) if the whole of derivatives are differentiable in the first form (1) :

$$[f^{(n)}(x, \alpha), g^{(n)}(x, \alpha)] + a_{n-1}(x)[f^{(n-1)}(x, \alpha), g^{(n-1)}(x, \alpha)] + \dots + a_0(x)[f(x, \alpha), g(x, \alpha)] = [g(x), g(x)]$$

(ii) if the whole of derivatives are differentiable in the second form (2) :

$$[g^{(n)}(x, \alpha), f^{(n)}(x, \alpha)] + a_{n-1}(x)[g^{(n-1)}(x, \alpha), f^{(n-1)}(x, \alpha)] + \dots + a_0(x)[g(x, \alpha), f(x, \alpha)] = [g(x), g(x)]$$

Then we have:

$$(10) \quad f^{(n)}(x, \alpha) + a_{n-1}(x)f^{(n-1)}(x, \alpha) + \dots + a_0(x)f(x, \alpha) = g(x)$$

$$(11) \quad g^{(n)}(x, \alpha) + a_{n-1}(x)g^{(n-1)}(x, \alpha) + \dots + a_0(x)g(x, \alpha) = g(x)$$

$$(12) \quad g^{(n)}(x, \alpha) + a_{n-1}(x)g^{(n-1)}(x, \alpha) + \dots + a_0(x)g(x, \alpha) = g(x)$$

$$(13) \quad f^{(n)}(x, \alpha) + a_{n-1}(x)f^{(n-1)}(x, \alpha) + \dots + a_0(x)f(x, \alpha) = g(x)$$

To obtain the Eqs. (10) and (13) and the Eqs.(11) and(12) are the same.

**Case 2:**  $a_i \leq 0, i = 0, \dots, n-1$

(i) if the whole of derivatives are differentiable in the first form (1) :

$$[f^{(n)}(x, \alpha), g^{(n)}(x, \alpha)] + a_{n-1}(x)[g^{(n-1)}(x, \alpha), f^{(n-1)}(x, \alpha)] + \dots + a_0(x)[g(x, \alpha), f(x, \alpha)] = [g(x), g(x)]$$

(ii) if the whole of derivatives are differentiable in the second form (2) :

$$[g^{(n)}(x, \alpha), f^{(n)}(x, \alpha)] + a_{n-1}(x)[f^{(n-1)}(x, \alpha), g^{(n-1)}(x, \alpha)] + \dots + a_0(x)[f(x, \alpha), g(x, \alpha)] = [g(x), g(x)]$$

$$\dots + a_0(x)[f(x, \alpha), g(x, \alpha) = [g(x), g(x)]]$$



Then we have:

$$(14) \quad f^{(n)}(x, \alpha) + a_{n-1}(x)g^{(n-1)}(x, \alpha) + \dots + a_0(x)g(x, \alpha) = g(x)$$

$$(15) \quad g^{(n)}(x, \alpha) + a_{n-1}(x)f^{(n-1)}(x, \alpha) + \dots + a_0(x)f(x, \alpha) = g(x)$$

$$(16) \quad g^{(n)}(x, \alpha) + a_{n-1}(x)f^{(n-1)}(x, \alpha) + \dots + a_0(x)f(x, \alpha) = g(x)$$

$$(17) \quad f^{(n)}(x, \alpha) + a_{n-1}(x)g^{(n-1)}(x, \alpha) + \dots + a_0(x)g(x, \alpha) = g(x)$$

To obtain the Eqs. ( 14) and ( 17) and the Eqs. (15) and (16) are the same.

**Case 3:**  $a_i \leq 0, i = 0, \dots, k, a_i \geq 0, i = k + 1, \dots, n - 1$

(i) if the whole of derivatives are differentiable in the first form (1) :

$$[f^{(n)}(x, \alpha), g^{(n)}(x, \alpha)] + a_{n-1}(x)[f^{(n-1)}(x, \alpha), g^{(n-1)}(x, \alpha)] + \dots + a_{k+1}(x)[f^{(k+1)}(x, \alpha), g^{(k+1)}(x, \alpha)] \\ + a_k(x)[g^{(k)}(x, \alpha), f^{(k)}(x, \alpha)] + \dots + a_0(x)[g(x, \alpha), f(x, \alpha)] = [g(x), g(x)]$$

(ii) if the whole of derivatives are differentiable in the second form (2) :

$$[g^{(n)}(x, \alpha), f^{(n)}(x, \alpha)] + a_{n-1}(x)[g^{(n-1)}(x, \alpha), f^{(n-1)}(x, \alpha)] + \dots + a_{k+1}(x)[g^{(k+1)}(x, \alpha), f^{(k+1)}(x, \alpha)] \\ + a_k(x)[f^{(k)}(x, \alpha), g^{(k)}(x, \alpha)] + \dots + a_0(x)[f(x, \alpha), g(x, \alpha)] = [g(x), g(x)]$$

Then we have:

$$(18) \quad f^{(n)}(x, \alpha) + a_{n-1}(x)f^{(n-1)}(x, \alpha) + \dots + a_{k+1}(x)f^{(k+1)}(x, \alpha) \\ + a_k(x)g^{(k)}(x, \alpha) + \dots + a_0(x)g(x, \alpha) = g(x)$$

$$(19) \quad g^{(n)}(x, \alpha) + a_{n-1}(x)g^{(n-1)}(x, \alpha) + \dots + a_{k+1}(x)g^{(k+1)}(x, \alpha) \\ + a_k(x)f^{(k)}(x, \alpha) + \dots + a_0(x)f(x, \alpha) = g(x)$$

$$(20) \quad g^{(n)}(x, \alpha) + a_{n-1}(x)g^{(n-1)}(x, \alpha) + \dots + a_{k+1}(x)g^{(k+1)}(x, \alpha) \\ + a_k(x)f^{(k)}(x, \alpha) + \dots + a_0(x)f(x, \alpha) = g(x)$$

$$(21) \quad f^{(n)}(x, \alpha) + a_{n-1}(x)f^{(n-1)}(x, \alpha) + \dots + a_{k+1}(x)f^{(k+1)}(x, \alpha) \\ + a_k(x)g^{(k)}(x, \alpha) + \dots + a_0(x)g(x, \alpha) = g(x)$$

To obtain the Eqs. (18) and (21) and the Eqs. (19) and (20) are equivalent.

#### 4 Solution of the generalizations of the differentiability of fuzzy functions by Adomian method

In this section we solve the fuzzy differential equations under generalized differentiability by Adomian decomposition method and restrictions of the method will be discussed.

We consider the Eqs. (3.1.1) and (3.1.2) , in section 3 and solve by Adomian decomposition method. Assume  $f^{\sim}, g^{\sim}$  are fuzzy functions of the independent crisp variables  $x$  and  $t$  in Eqs. (3.1.1) and (3.1.2)

We have:

$$f^{\sim(n)}(x, \alpha) + a_{n-1}(x)f^{\sim(n-1)}(x, \alpha) + \dots + a_0(x)f^{\sim}(x, \alpha) = g(x)$$

$$g^{\sim(n)}(x, \alpha) + a_{n-1}(x)g^{\sim(n-1)}(x, \alpha) + \dots + a_0(x)g^{\sim}(x, \alpha) = g(x)$$

Definition:

$$(22) \quad g_1 \sim = g \sim, g_2 \sim = g \sim', \dots, g_n \sim = g \sim^{(n-1)} \quad \text{and} \quad f_1 \sim = f \sim, f_2 \sim = f \sim', \dots, f_n \sim = f \sim^{(n-1)}$$

$$(23) \quad f_i(0) = \gamma_{i1}(\alpha), \quad \text{and} \quad g_i(0) = \gamma_{i2}(0), \quad i = 1, 2, \dots, n$$

For solving this equation by Adomian decomposition method the equation should be in canonical form which can be derived by rewriting Eq. (22) as follows:

$$(24) \quad Lf_1 \sim = f_2 \sim, Lf_2 \sim = f_3 \sim, \dots, Lf_n \sim = g(x) - a_{n-1}(x)f_n \sim - \dots - a_0(x)f_1 \sim$$

$$(25) \quad Lg_1 \sim = g_2 \sim, Lg_2 \sim = g_3 \sim, \dots, Lg_n \sim = g(x) - a_{n-1}(x)g_n \sim - \dots - a_0(x)g_1 \sim$$

where  $L_t = \frac{\partial}{\partial t}$ , with the inverse operator  $L_t^{-1} = \int_0^t (\cdot) dt$ .

Applying the inverse operator, we get

$$(26) \quad f_i \sim = f_i(0) + \int_0^x f_{i+1} \sim(x) dx, \quad i = 1, \dots, n-1$$

$$(27) \quad f_n \sim = f_n(0) + \int_0^x (g(x) - a_{n-1}(x)f_n \sim - \dots - a_0(x)f_1 \sim) dx$$

$$(28) \quad g_i \sim = g_i(0) + \int_0^x g_{i+1} \sim(x) dx,$$

$$(29) \quad g_n \sim = g_n(0) + \int_0^x (g(x) - a_{n-1}(x)g_n \sim - \dots - a_0(x)g_1 \sim) dx$$

Adomian decomposition method considers the solutions  $f \sim, g \sim$  as the sum of a series as:

$$(30) \quad f_i \sim = \sum_{j=0}^{\infty} f_{i,j} \sim, \quad g_j \sim = \sum_{j=0}^{\infty} g_{i,j} \sim, \quad i = 1, \dots, n$$

So we can calculate the terms of  $f_i^{\sim} = \sum_{j=0}^{\infty} f_{i,j}^{\sim}$ ,  $g_j^{\sim} = \sum_{j=0}^{\infty} g_{i,j}^{\sim}$  term by term as long as we derive desired accuracy, the more terms the more accuracy. Therefore we have :

$$f^{\sim}_{i,0} + f^{\sim}_{i,1} + \dots = \gamma_{i1}(\alpha) + \int_0^x (f^{\sim}_{i+1,0} + f^{\sim}_{i+1,1} + \dots) dx,$$

$$i = 1, \dots, n-1$$

$$f^{\sim}_{n,0} + f^{\sim}_{n,1} + \dots = \gamma_{n1}(\alpha) + \int_0^x g(x) - a_{n-1}(x)(f^{\sim}_{n,0} + f^{\sim}_{n,1} + \dots) - \dots - a_0(x)(f^{\sim}_{1,0} + f^{\sim}_{1,1} + \dots) dx.$$

$$(31) \quad g^{\sim}_{i,0} + g^{\sim}_{i,1} + \dots = \gamma_{i2}(\alpha) + \int_0^x (g^{\sim}_{i+1,0} + g^{\sim}_{i+1,1} + \dots) dx, \quad i = 1, \dots, n-1$$

$$(32) \quad g^{\sim}_{n,0} + g^{\sim}_{n,1} + \dots = \gamma_{n2}(\alpha) + \int_0^x g(x) - a_{n-1}(x)(g^{\sim}_{n,0} + g^{\sim}_{n,1} + \dots) - \dots - a_0(x)(g^{\sim}_{1,0} + g^{\sim}_{1,1} + \dots) dx.$$

By solving (31) and (32) we can calculate the terms of series above.

Two other case are the same.

### 5 Examples

**Example 1.** Consider the following fuzzy equation with the indicated initial conditions:

$$(33) \quad y''' + y'' + y = 0, \quad y(0) = 0^{\sim}, \quad y'(0) = 1^{\sim}, \quad y''(0) = 2^{\sim}.$$

The exact solution are  $f^{\sim} = \gamma_{i1}(\alpha)e^x$  and  $g^{\sim} = \gamma_{i2}(\alpha)e^x$  for  $\alpha \in [0,1]$ .

Since coefficients are positive then we have case 1. of section 3.

$$(34) \quad f'''(x, \alpha) + f''(x, \alpha) + f(x, \alpha) = 0, \quad g'''(x, \alpha) + g''(x, \alpha) + g(x, \alpha) = 0,$$

By Adomian decomposition method we have:

$$(35) \quad f^{\sim}_1 = f^{\sim}, \quad f^{\sim}_2 = f^{\sim \prime}, \quad f^{\sim}_3 = f^{\sim \prime \prime},$$

and

$$g^{\sim}_1 = g^{\sim}, \quad g^{\sim}_2 = g^{\sim \prime}, \quad g^{\sim}_3 = g^{\sim \prime \prime},$$

Then:

$$(36) \quad Lf^{\sim}_1 = f^{\sim}_2, \quad Lf^{\sim}_2 = f^{\sim}_3, \quad Lf^{\sim}_3 = -f^{\sim}_3 - f^{\sim}_1$$

$$Lg_1 = g_2, \quad Lg_2 = g_3, \quad Lg_3 = -g_3 - g_1,$$

Then we have:

$$(37) \quad f^{\sim}_1 = (0 - \alpha) + \int_0^x f^{\sim}_2 dx,$$

$$f^{\sim}_2 = (1 - \alpha) + \int_0^x f^{\sim}_3 dx,$$

$$f^{\sim}_3 = (2 - \alpha) - \int_0^x (f^{\sim}_3 + f^{\sim}_1) dx.$$

$$(38) \quad \begin{aligned} g_{\sim 1} &= (0 + \alpha) + \int_0^x g_{\sim 2} dx, \\ g_{\sim 2} &= (1 + \alpha) + \int_0^x g_{\sim 3} dx, \\ g_{\sim 3} &= (2 + \alpha) - \int_0^x (g_{\sim 3} + g_{\sim 1}) dx. \end{aligned}$$

Considers the sum of series as:

$$f_i^{\sim} = \sum_{j=0}^{\infty} f_{i,j}^{\sim}, g_j^{\sim} = \sum_{i=0}^{\infty} g_{i,j}^{\sim}, i=1,2,3$$

Then

$$(39) \quad \begin{aligned} f_{\sim 2,0} + f_{\sim 2,1} + \dots &= 1 - \alpha + \int_0^x (f_{\sim 3,0} + f_{\sim 3,1} + \dots) dx, \\ f_{\sim 1,0} + f_{\sim 1,1} + \dots &= -\alpha + \int_0^x (f_{\sim 2,0} + f_{\sim 2,1} + \dots) dx, \\ f_{\sim 3,0} + f_{\sim 3,1} + \dots &= 2 - \alpha - \int_0^x (f_{\sim 3,0} + f_{\sim 3,1} + \dots + f_{\sim 1,0} + f_{\sim 1,1} + \dots) dx, \end{aligned}$$

$$(40) \quad \begin{aligned} g_{\sim 1,0} + g_{\sim 1,1} + \dots &= \alpha + \int_0^x (g_{\sim 2,0} + g_{\sim 2,1} + \dots) dx, \\ g_{\sim 2,0} + g_{\sim 2,1} + \dots &= 1 + \alpha + \int_0^x (g_{\sim 3,0} + g_{\sim 3,1} + \dots) dx, \\ g_{\sim 3,0} + g_{\sim 3,1} + \dots &= 2 + \alpha - \int_0^x (g_{\sim 3,0} + g_{\sim 3,1} + \dots + g_{\sim 1,0} + g_{\sim 1,1} + \dots) dx, \end{aligned}$$

Then we obtain

$$(41) \quad f_{\sim 1,0} = -\alpha, f_{\sim 2,0} = 1 - \alpha, f_{\sim 3,0} = 2 - \alpha, \text{ and}$$

$$g_{1,0} \sim \alpha, \quad g_{2,0} \sim 1 + \alpha, \quad g_{3,0} \sim 2 + \alpha,$$

(42)

$$f_{1,n+1} \sim \int_0^x f_{2,n} \sim dx,$$

$$f_{2,n+1} \sim \int_0^x f_{3,n} \sim dx,$$

$$f_{3,n+1} \sim -\int_0^x (f_{3,n} + f_{1,n}) \sim dx, \quad n = 0, 1, \dots$$

$$g_{1,n+1} \sim \int_0^x g_{2,n} \sim dx,$$

$$g_{3,n+1} \sim -\int_0^x (g_{3,n} + g_{1,n}) \sim dx, \quad n = 0, 1, \dots \quad g_{2,n+1} \sim \int_0^x g_{3,n} \sim dx,$$

For  $n = 0$  we obtain:

(43)

$$f_{1,1} \sim \int_0^x f_{2,0} \sim dx = \int_0^x (1 - \alpha) \sim dx = (1 - \alpha)x,$$

$$f_{2,1} \sim \int_0^x f_{3,0} \sim dx = \int_0^x (2 - \alpha) \sim dx = (2 - \alpha)x,$$

$$f_{3,1} \sim -\int_0^x (f_{3,0} + f_{1,0}) \sim dx = -\int_0^x (2 - \alpha - \alpha) \sim dx = -(2 - 2\alpha)x,$$

(44)

$$g_{1,1} \sim \int_0^x g_{2,0} \sim dx = \int_0^x (1 + \alpha) \sim dx = (1 + \alpha)x,$$

$$g_{2,1} \sim \int_0^x g_{3,0} \sim dx = \int_0^x (2 + \alpha) \sim dx = (2 + \alpha)x,$$

$$g_{3,1}^{\sim} = -\int_0^x (g_{3,0}^{\sim} + g_{1,0}^{\sim}) dx = -\int_0^x (2 + \alpha + \alpha) dx = -(2 + 2\alpha)x,$$

For  $n = 1$  we obtain:

$$(45) \quad f_{1,2}^{\sim} = \int_0^x f_{2,1}^{\sim} dx = \frac{2-\alpha}{2} x^2,$$

$$f_{2,2}^{\sim} = \int_0^x f_{3,1}^{\sim} dx = \frac{-(2-2\alpha)}{2} x^2,$$

$$f_{3,2}^{\sim} = \frac{1-\alpha}{2} x^2,$$

$$(46) \quad g_{1,2}^{\sim} = \frac{2+\alpha}{2} x^2,$$

$$g_{2,2}^{\sim} = \frac{-(2+2\alpha)}{2} x^2,$$

$$g_{3,2}^{\sim} = \frac{1+\alpha}{2} x^2,$$

Since  $f_1^{\sim} = f^{\sim}$ ,  $g_1^{\sim} = g^{\sim}$  and according to generalized differentiability  $[y(t)]_{\alpha} = [f(t, \alpha), g(t, \alpha)] = [f^{\sim}, g^{\sim}]$  we have:

$$(47) \quad f^{\sim} = f_1^{\sim} = \sum_{j=0}^{\infty} f_{1,j}^{\sim} = f_{1,0}^{\sim} + f_{1,1}^{\sim} + \dots = -\alpha + (1-\alpha)x + \frac{2-\alpha}{2} x^2 + \dots$$

$$g^{\sim} = g_1^{\sim} = \sum_{j=0}^{\infty} g_{1,j}^{\sim} = g_{1,0}^{\sim} + g_{1,1}^{\sim} + \dots = \alpha + (1+\alpha)x + \frac{2+\alpha}{2} x^2 + \dots$$

In this example we have derived the exact solution.

**Example 2.** Consider the following fuzzy equation:

$$(48) \quad y'' - y' + y = x,$$



$$y(0) = 0^{\sim}, y'(0) = 1^{\sim}$$

Solve the fuzzy differential equation under generalized differentiability by apply Adomian decomposition method.

We have:

$$(49) \quad \begin{aligned} f'' - g' + f &= x, \\ g'' - f' + g &= x, \\ f^{\sim}_1 &= f, f^{\sim}_2 = f' \text{ and } g^{\sim}_1 = g, g^{\sim}_2 = g', \end{aligned}$$

By apply Adomian decomposition method on above equations, we have:

$$(50) \quad f^{\sim}_{1,0} = -\alpha, f^{\sim}_{2,0} = 1 - \alpha, g^{\sim}_{1,0} = \alpha, g^{\sim}_{2,0} = 1 + \alpha,$$

$$(51) \quad f^{\sim}_{1,n+1} = \int_0^x f^{\sim}_{2,n} dx, f^{\sim}_{2,n+1} = \int_0^x (x + g^{\sim}_{2,n} - f^{\sim}_{1,n}) dx,$$

$$g^{\sim}_{1,n+1} = \int_0^x g^{\sim}_{2,n} dx, g^{\sim}_{2,n+1} = \int_0^x (x + f^{\sim}_{2,n} - g^{\sim}_{1,n}) dx, n = 0,1,2,\dots$$

We obtain:

$$\begin{aligned} f^{\sim}_1 &= f^{\sim} = f^{\sim}_{1,0} + f^{\sim}_{1,1} + f^{\sim}_{1,2} + \dots \\ &= x + \frac{x^2}{2} - \frac{x^3}{6} - \alpha(1+x) + \dots \end{aligned}$$

$$(52) \quad \begin{aligned} g^{\sim}_1 &= g^{\sim} = g^{\sim}_{1,0} + g^{\sim}_{1,1} + g^{\sim}_{1,2} + \dots \\ &= x + \frac{1-2\alpha}{2} x^2 + \frac{x^3}{6} + \alpha(1+x) + \dots \end{aligned}$$

## **6 Conclusions and future research**

In this paper we presented the generalized differentiability for the n-th order, linear, differential equation having fuzzy initial conditions. We apply Adomian decomposition method.

This method has a useful feature in that it provides the solution in a rapid convergent power series with elegantly computable convergence of the solution.

So, future research will be concerned with fuzzy  $a_i$ , and fuzzy initial conditions.

## References

- [1] Zadeh, L., Toward a generalized theory of uncertainty (GTU)-an outline, *Information Sciences* 172, 1-40, 2005.
- [2] Bede, B., and Gal, S.G., Almost periodic fuzzy- number-valued functions, *Fuzzy* 147, 385-403, 2004.
- [3] ] Bede, B., and Gal, S. G., Generalizations of the differentiability of fuzzy- with applications to fuzzy differential equations, *Fuzzy* 151, 581-599, 2005.
- [4] Dubois, D., and Prade, H., Fuzzy numbers: an overview, in: J. Bezdek (Ed.), *Analysis of Fuzzy Information*, CRC Press, 112-148, 1987.
- [5] Wu, C., and Gong, Z., On Henstock .integral of fuzzy number-valued functions, I, *Fuzzy sets and systems* 120, 523-532, 2001.
- [6] Gal, S. G., Approximation theory in fuzzy setting , in: G.A. Anastassiou (Ed.), *Handbook of Analytic. Computational Methods in Applied Mathematics*, Chapman & Hall/CRC press, 617-666, 2000.
- [7] Chalco-Cano, Y., Roman-Flores, on new solutions of fuzzy differential equations, *Chaos Solutions and Fractals*, In press, 2006.
- [8] Puri, M., and Ralescu, D., Differentials of fuzzy functions, *Journal of Mathematical Analysis and Applications* 91, 552-558, 1983.