



Well- posedness of the Rothe difference scheme for reverse parabolic equations

Allaberen Ashyralyev^a, Ayfer Dural^b, Yaşar Sözen^{c*}

^aDepartment of Mathematics, Fatih University, Istanbul, 34500, Turkey

^bGaziosman Paşa Lisesi Istanbul, Turkey

Abstract

We consider the Rothe difference scheme for approximate solution of the abstract parabolic equation in a Hilbert space with the nonlocal boundary condition. Theorems on stability estimates, coercivity and almost coercivity estimates for the solution of this difference scheme are established. In application, new coercivity inequalities for the solution of multi-point nonlocal boundary value difference equations of parabolic type are obtained.

Keywords: Multipoint nonlocal boundary value problem, parabolic equations, reverse type, difference equations, well-posedness, almost coercivity 2000 MSC: 47D06, 35K20

* Corresponding author. E-mail: ysozen@fatih.edu.tr

1 Introduction

In the study of boundary value problems for partial differential equations, the role played by, well-posedness (coercivity inequality) is well known (see Ladyzhenskaya et al. 1968, Ladyzhenskaya and Ural'tseva 1968, Vishik et al. 1959). Coercivity inequalities for nonlocal boundary value problems for partial differential equations parabolic and elliptic types have been studied extensively by many researchers see Aibeche and Favini (2005), Clement and Guerre (1999), Shakhmurov (2004), Sobolevskii (1971) and references given therein.

In the paper (see Ashyralyev and el al. 2008) we considered the abstract nonlocal boundary value problem

$$\begin{cases} \frac{du(t)}{dt} - Au(t) = f(t) & (0 \leq t \leq 1), \\ u(1) = \sum_{k=1}^p \alpha_k u(\theta_k) + \varphi, \\ 0 \leq \theta_1 < \theta_2 < \dots < \theta_p < 1 \end{cases} \quad (1)$$

in a Hilbert space H with self-adjoint positive definite operator A , under the assumption

$$\sum_{k=1}^p |\alpha_k| \leq 1 \quad (2)$$

The well-posedness of multi-point nonlocal boundary value problem (1) in spaces $C_1^\alpha(H)$ and $C^\alpha(H)$ was established. Moreover, as applications, these abstract results enabled us to obtain new coercivity estimates in various Hölder norms for the solutions of nonlocal boundary value problems for parabolic equations.

In the present article, our focus is the well-posedness of the first order of accuracy Rothe difference scheme

$$\begin{cases} \tau^{-1}(u_k - u_{k-1}) - Au_{k-1} = \varphi_k, \varphi_k = f(t_k), \\ t_k = k\tau, 1 \leq k \leq N, N\tau = 1, \\ u_N = \sum_{m=1}^p \alpha_m u_{\ell_m} + \varphi, \\ \ell_m = \left[\begin{matrix} \theta_m \\ \tau \end{matrix} \right], 1 \leq m \leq p \end{cases} \quad (3)$$

for approximately solving problem (1).

Let $[0,1]_\tau = \{t_k = k\tau, k = 1, \dots, N, N\tau = 1\}$ be the *uniform grid space* with step size $\tau > 0$, where N is a fixed positive integer.

Throughout the paper, $F([0,1]_\tau, H)$ denotes the linear space of grid functions $\varphi^\tau = \{\varphi_k\}_1^N$ with values in the Hilbert space H.

Let $C_\tau(H) = C([0,1]_\tau, H)$ be the Banach space of bounded grid functions with the norm

$$\begin{aligned} \|\varphi^\tau\|_{C^\alpha(H)} &= \|\varphi^\tau\|_{C(H)} + \max_{1 \leq k < k+r \leq N} \frac{\|\varphi_{k+r} - \varphi_k\|_H}{(r\tau)^\alpha}, \\ \|\varphi^\tau\|_{C^\alpha(H)} &= \|\varphi^\tau\|_{C(H)} + \max_{1 \leq k < k+r \leq N} \frac{((N-k)\tau)^\alpha \|\varphi_{k+r} - \varphi_k\|_H}{(r\tau)^\alpha}. \end{aligned}$$

We say that difference problem (3) is *stable* in $F([0,1]_\tau, H)$, if we have the following *stability estimate*

$$\|\{u_{k-1}\}_1^N\|_{F([0,1]_\tau, H)} \leq M \left(\|\varphi^\tau\|_{F([0,1]_\tau, H)} + \|\varphi\|_H \right),$$

where M is independent of φ^τ, φ and τ .

Difference problem (3) is said to be *well-posed* in $F([0,1]_\tau, H)$, if for every

$\varphi^\tau \in F([0, 1]_\tau, H)$ problem (3) is uniquely solvable and have the following *coercivity estimate* :

$$\begin{aligned} & \left\| \left\{ \tau^{-1} (u_k - u_{k-1}) \right\}_1^N \right\|_{F([0,1]_\tau, H)} + \left\| \{ Au_{k-1} \}_1^N \right\|_{F([0,1]_\tau, H)} \\ & \leq M \left(\left\| \phi^\tau \right\|_{F([0,1]_\tau, H)} + \|A\phi\|_{H'} \right), \end{aligned}$$

where $H' \subset H$, M does not depend on φ^τ, φ and τ .

Throughout the paper, M shall indicate positive constants which can be different from time to time and we are not interested to precise. We shall write $M(\alpha, \beta, \dots)$ to stress the fact that the constant depends only on α, β, \dots .

2 The First Order of Accuracy Difference Scheme

Let us start with some auxiliary lemmas we need below. Throughout the paper, H denotes a Hilbert space and A is a positive definite self-adjoint operator with $A \geq \delta I$ for some $\delta > 0$.

Lemma 1.1. (See Ashyralyev and Sobolevskii (1994)). The following estimates hold:

$$\left\| t^k A^k e^{-tA} \right\|_{H \rightarrow H} \leq M, \quad t > 0, \quad K \geq 0, \quad (4)$$

$$\left\| R^k \right\|_{H \rightarrow H} \leq \frac{1}{(1 + \delta\tau)^k}, \quad K \geq 1, \quad (5)$$

$$\left\| \tau A R^k \right\|_{H \rightarrow H} \leq \frac{1}{k}, \quad K \geq 1, \quad (6)$$

$$\left\| A^\beta (R^{k+r} - R^k) \right\|_{H \rightarrow H} \leq M \frac{(r\tau)^\gamma}{(k\tau)^{\beta+\gamma}}, \quad 1 \leq k < k+r \leq N, \beta \in \{0,1\}, 0 \leq \gamma \leq 1, \quad (7)$$

for some $M, \delta > 0$, which are independent of τ is a positive small number and

$R = (I + \tau A)^{-1}$ is the resolvent of A .

Lemma 1.2. Assume that (2) holds. Then, the operator

$$I - \sum_{k=1}^p \alpha_k R^{N \cdot \left[\frac{\theta_k}{r} \right]} \tag{8}$$

has an inverse

$$T_\tau = \left(I - \sum_{k=1}^p \alpha_k R^{N \cdot \left[\frac{\theta_k}{r} \right]} \right)^{-1}$$

and the following estimate is satisfied:

$$\|T_\tau\|_{H \rightarrow H} \leq C(\delta, \theta_p). \tag{9}$$

Proof. The proof of estimate (9) is based on the triangle inequality, assumption (2), and the estimate

$$\left\| \left(I - \sum_{k=1}^p \alpha_k R^{N \cdot \left[\frac{\theta_k}{\tau} \right]} \right)^{-1} \right\|_{H \rightarrow H} \leq \sup_{\delta \leq \mu} \frac{1}{\left| 1 - \sum_{k=1}^p \alpha_k (1 + \tau\mu)^{-N \cdot \left[\frac{\theta_p}{\tau} \right]} \right|}.$$

□

Let us now obtain the formula for the solution of problem (3). It is clear that the first order of accuracy difference scheme

$$\begin{cases} \tau^{-1}(u_k - u_{k-1}) - Au_{k-1} = \varphi_k, \varphi_k = f(t_k), \\ t_k = k\tau, 1 \leq k \leq N, N\tau = 1, \\ u_N = \sum_{m=1}^p \alpha_m u_{\ell_m} + \varphi, \\ \ell_m = \left[\frac{\theta_m}{\tau} \right], 1 \leq m \leq p \end{cases} \tag{10}$$

has a solution and the following formula holds:

$$u_k = R^{N-k} u_N - \sum_{j=k+1}^N R^{j-k} \varphi_j \tau, \quad 0 \leq k \leq N-1. \quad (11)$$

Applying formula (11) and the nonlocal boundary condition

$$\xi = u_N = \sum_{m=1}^p \alpha_m u_{\ell_m} + \varphi,$$

we can write

$$\xi = \sum_{k=1}^p \alpha_k \left(R^{N-\ell_k} \xi - \sum_{j=\ell_k+1}^N R^{j-\ell_k} \varphi_j \tau \right) + \varphi.$$

Using Lemma 1.2, we get

$$u_N = T_\tau \left(- \sum_{k=1}^p \sum_{j=\ell_k+1}^N \alpha_k R^{j-\ell_k} \varphi_j \tau + \varphi \right) \quad (12)$$

Hence, difference equation (10) is uniquely solvable and for the solution, formulas (11) and (12) are valid.

Theorem 1.3. Suppose that (2) holds and $\varphi \in D(A)$. Then, for the solution of difference scheme (10) the following stability estimate

$$\max_{0 \leq k \leq N} \|u_k\|_H \leq C(\delta, \theta_p) \left(\|\varphi\|_H + \|\varphi^\tau\|_{C_\tau(H)} \right). \quad (13)$$

holds, where $C(\delta, \theta_p)$ is independent of τ, φ , and φ^τ .

Proof. From estimate (5), formula (11), and $N\tau = 1$ it follows that

$$\max_{0 \leq k \leq N-1} \|u_k\|_H \leq \|u_N\|_H + \max_{1 \leq j \leq N} \|\varphi_j\|_H.$$

Using assumption (2), estimate (5), (9), formula (12), and $N_\tau = 1$, we obtain

$$\|u_N\|_H \leq C_1(\delta, \theta_p) \left(\|\varphi\|_H + \|\varphi^\tau\|_{C_\tau(H)} \right)$$

From these estimates it follows (13).

This concludes the proof of Theorem 1.3. □

It is well-known that problem (1) in the space $C([0,1], H)$ is not well-posed for the general positive definite self-adjoint operator A and Hilbert space H . Hence, the well-posedness of difference problem (10) in $C([0,1]_\tau, H)$ norm does not take place uniformly with respect to $\tau > 0$.

Theorem 1.4. Let (2) holds and $\varphi \in D(A)$. then, for the solution of difference problem (10), the almost coercivity inequality

$$\begin{aligned} & \left\| \left\{ \tau^{-1}(u_k - u_{k-1}) \right\}_1^N \right\|_{C_\tau(H)} + \left\| \left\{ Au_{k-1} \right\}_1^N \right\|_{C_\tau(H)} \\ & \leq C(\delta, \theta_p) \left(\min \left\{ \ln \frac{1}{\tau}, 1 + \ln \|A\|_{H \rightarrow H} \right\} \cdot \|\varphi^\tau\|_{C_\tau(H)} + \|A\varphi\|_H \right) \end{aligned} \tag{14}$$

is valid, where $C(\delta, \theta_p)$ does not depend on τ, φ , and φ^τ .

Proof. Using formula (11), estimate (5), we get for $1 \leq k \leq N$

$$\|Au_{k-1}\|_H \leq \|Au_N\|_H + \|\varphi^\tau\|_{C_\tau(H)} \sum_{j=k}^N \|\tau AR^{j-k+1}\|_{H \rightarrow H}. \tag{15}$$

It follows from Theorem 1.2 (see Ashyralyev and Sobolevskii (1994) on page 87) that

$$\begin{aligned} \sum_{j=k}^N \|\tau AR^{j-k+1}\|_{H \rightarrow H} &= \sum_{m=1}^{N-k+1} \tau \|AR^m\|_{H \rightarrow H} \\ &\leq M \min \left\{ \ln \frac{1}{\tau}, 1 + |\ln \|A\|_{H \rightarrow H}| \right\}. \end{aligned} \tag{16}$$

By formula (12), estimate (9), and assumption (2), we obtain

$$\|Au_N\|_H \leq C(\delta, \theta_p) \left(\|\varphi^\tau\|_{C_\tau(H)} \min \left\{ \ln \frac{1}{\tau}, |\ln \|A\|_{H \rightarrow H}| \right\} + \|A\varphi\|_H \right). \tag{17}$$

Thus, from estimates (15)- (17) it follows that

$$\begin{aligned} & \left\| \{Au_{k-1}\}_1^N \right\|_{C_\tau(H)} \leq C(\delta, \theta_p) \\ & \times \left(\left\| \varphi^\tau \right\|_{C_\tau(H)} \min \left\{ \ln \frac{1}{\tau}, \left| \ln \|A\|_{H \rightarrow H} \right| \right\} + \|A\varphi\|_H \right). \end{aligned} \quad (18)$$

Using difference equation (10), the triangle inequality, and estimate (18), we get estimate (14).

This completes the proof of Theorem 1. \square

Theorem 1.5. Suppose that (2) holds and $\varphi \in D(A)$. Then, the solution of difference scheme (10) satisfy the following stability estimate

$$\begin{aligned} & \left\| \{\tau^{-1}(u_k - u_{k-1})\}_1^N \right\|_{C_\tau^\alpha(H)} + \left\| \{Au_{k-1}\}_1^N \right\|_{C_1^\alpha(H)} \\ & \leq C(\delta, \theta_p) \left(\frac{1}{\alpha(1-\alpha)} \cdot \left\| \varphi^\tau \right\|_{C_1^\alpha(H)} + \|A\varphi\|_H \right), \end{aligned} \quad (19)$$

where $C(\delta, \theta_p)$ is independent of τ, φ , and φ^τ .

Proof. It follows from formula (11) and identity

$$\tau AR = I - R \quad (20)$$

that for $1 \leq k \leq N$

$$Au_{k-1} = R^{N-k+1} Au_N - \sum_{j=k}^N \tau AR^{j-k+1} (\varphi_j - \varphi_{k-1}) + (R^{N-k+1} - I) \varphi_{k-1}. \quad (21)$$

Thus, using estimate (5), (6), and the definition of $C_1^\alpha(H)$ -norm, we get for $1 \leq k \leq N$

$$\begin{aligned} \|Au_{k-1}\|_H &\leq \|Au_N\|_H \\ &+ \frac{\|\varphi^\tau\|_{C_1^\alpha(H)}}{((N-k+1)\tau)^\alpha} \sum_{j=k}^N \frac{\tau}{((j-k+1)\tau)^{1-\alpha}} + 2\|\varphi^\tau\|_{C_1^\alpha(H)} \quad (22) \\ &\leq \|Au_N\|_H + \frac{4}{\alpha} \|\varphi^\tau\|_{C_1^\alpha(H)}. \end{aligned}$$

Now, we estimate $\|Au_N\|_H$.

From formula (12) and $\tau AR = I - R$ it follows that

$$Au_N = T_\tau \left\{ - \sum_{k=1}^p \alpha_k \left(\sum_{j=\ell_k+1}^N \tau AR^{j-\ell_k} (\varphi_j - \varphi_{\ell_k}) + (I - R^{N-\ell_k}) \varphi_{\ell_k} \right) + A\varphi \right\}.$$

Hence, by estimates (5), (6), (9), the definition of $C_1^\alpha(H)$ -norm, and assumption (2), we obtain

$$\|Au_N\|_H \leq C(\delta, \theta_p) \left(\frac{4}{\alpha} \|\varphi^\tau\|_{C_1^\alpha(H)} + \|A\varphi\|_H \right). \quad (23)$$

Thus, from estimates (22), (23) it follows that

$$\left\| \{Au_{k-1}\}_1^N \right\|_{C_\tau(H)} \leq C(\delta, \theta_p) \left(\frac{1}{\alpha} \|\varphi^\tau\|_{C_1^\alpha(H)} + \|A\varphi\|_H \right). \quad (24)$$

Let us now estimate

$$\max_{1 \leq k < k+r \leq N} \frac{((N-k+1)\tau)^\alpha \|Au_{k-1+r} - Au_{k-1}\|_H}{(r\tau)^\alpha}.$$

First, let $N - k + r \leq 2r$. By estimate (24) and the triangle inequality, we obtain

$$\frac{((N-k+1)\tau)^\alpha \|Au_{k-1+r} - Au_{k-1}\|_H}{(r\tau)^\alpha} \leq C(\delta, \theta_p) \left(\frac{1}{\alpha} \|\varphi^\tau\|_{C_1^\alpha(H)} + \|A\varphi\|_H \right). \quad (25)$$

Next, let $N - k + r \leq 2r$. From formula (11) it follows that

$$\begin{aligned}
 Au_{k-1} - Au_{k-1+r} &= (R^{N-k+1} - R^{N-k+1-r}) Au_N \\
 &\quad - \sum_{j=k}^{k+2r-2} \tau AR^{j-k+1} (\phi_j - \phi_{k-1}) \\
 &\quad + \sum_{j=k+r}^{k+2r-2} \tau AR^{j-(k-1+r)} (\phi_j - \phi_{k-1+r}) \\
 &\quad - \sum_{j=k+2r-1}^N \tau A (R^{j-k+1} - R^{j-(k-1+r)}) (\phi_j - \phi_{k-1}) \\
 &\quad + (I - R^{r-1}) (\phi_{k-1+r} - \phi_{k-1}) + (R^{N-k+1} - R^{N-(k-1+r)}) \phi_{k-1} \\
 &= I_1(k) + I_2(k) + I_3(k) + I_4(k) + I_5(k) + I_6(k).
 \end{aligned} \tag{26}$$

We first estimate $I_1(k)$. Using estimates (7) for $\beta = 0$, and the fact $N - j + 1 > 2r$, we get

$$\|I_1(k)\|_H \leq C(\delta, \theta_p) \frac{(r\tau)^\alpha}{((N - k + 1)\tau)^\alpha} \left(\frac{1}{\alpha} \|\varphi^\tau\|_{C_1^\alpha(H)} + \|A\varphi\|_H \right). \tag{27}$$

Next, it follows from estimate (6) and the definition of $C_1^\alpha(H)$ -norm that

$$\begin{aligned}
 \|I_2(k)\|_H &\leq \frac{\|\varphi^\tau\|_{C_1^\alpha(H)}}{((N - k + 1)\tau)^\alpha} \sum_{j=k+r}^{k+2r-2} \frac{\tau}{((j - (k - 1 + r))\tau)^{1-\alpha}} \\
 &\leq \frac{2^\alpha}{\alpha} \frac{(r\tau)^\alpha}{((N - k + 1)\tau)^\alpha} \|\varphi^\tau\|_{C_1^\alpha(H)}.
 \end{aligned} \tag{28}$$

By using estimate (6), the definition of $C_1^\alpha(H)$ -norm, and the fact $N - j + 1 > 2r$, we obtain

$$\begin{aligned}
 \|I_3(k)\|_H &\leq \frac{2^\alpha \|\varphi^\tau\|_{C_1^\alpha(H)}}{((N - k + 1)\tau)^\alpha} \sum_{j=k+r}^{k+2r-2} \frac{\tau}{((j - (k - 1 + r))\tau)^{1-\alpha}} \\
 &\leq \frac{2^\alpha \|\varphi^\tau\|_{C_1^\alpha(H)}}{((N - k + 1)\tau)^\alpha} \frac{(r\tau)^\alpha}{\alpha}.
 \end{aligned} \tag{29}$$

If follows from estimate (7) for $\beta = 1$, the definition of $C_1^\alpha(H)$ -norm, and the fact $j - k + 1 > 2r$, that

$$\begin{aligned} \|I_4(k)\|_H &\leq M \frac{2^\alpha \|\varphi^\tau\|_{C_1^\alpha(H)}}{((N - k + 1)\tau)^\alpha} r\tau \sum_{j=k+2r-1}^N \frac{\tau}{((j - (k - 1 + r))\tau)^{2-\alpha}} \\ &\leq M \frac{2^\alpha \|\varphi^\tau\|_{C_1^\alpha(H)}}{((N - k + 1)\tau)^\alpha} \frac{(r\tau)^\alpha}{(1 - \alpha)}. \end{aligned} \quad (30)$$

Using estimate (5), the definition of $C_1^\alpha(H)$ -norm, we obtain

$$\|I_5(k)\|_H \leq \frac{2^\alpha \|\varphi^\tau\|_{C_1^\alpha(H)}}{((N - k + 1)\tau)^\alpha} (r\tau)^\alpha. \quad (31)$$

Finally, from estimate (7) for $\beta = 0$ and the fact $N - j + 1 > 2r$ it follows that

$$\|I_6(k)\|_H \leq 2^\alpha \|\varphi^\tau\|_{C_1^\alpha(H)}. \quad (32)$$

Thus, combining estimates (27)-(32), we get for $N - j + 1 > 2r$

$$\frac{((N - k + 1)\tau)^\alpha \|Au_{k-1+r} - Au_{k-1}\|_H}{(r\tau)^\alpha} \leq C(\delta, \theta_p) \left(\frac{\|\varphi^\tau\|_{C_1^\alpha(H)}}{\alpha(1 - \alpha)} + \|A\varphi\|_H \right). \quad (33)$$

From estimates (25) and (33) it follows that

$$\max_{1 \leq k < k+r \leq N} \frac{((N - k + 1)\tau)^\alpha \|Au_{k-1+r} - Au_{k-1}\|_H}{(r\tau)^\alpha} \leq C(\delta, \theta_p) \left(\frac{\|\varphi^\tau\|_{C_1^\alpha(H)}}{\alpha(1 - \alpha)} + \|A\varphi\|_H \right). \quad (34)$$

Combining estimates (24)-(34), we obtain that

$$\left\| \{Au_{k-1}\}_1^N \right\|_{C\tau(H)} \leq C(\delta, \theta_p) \left(\frac{1}{\alpha(1 - \alpha)} \|\varphi^\tau\|_{C_1^\alpha(H)} + \|A\varphi\|_H \right). \quad (35)$$

Hence, estimate (19) follows from difference equation (10), estimate (35) and the triangle inequality.

This concludes the proof of Theorem 1.5. □

Let $H_\alpha = H_{\alpha,\infty}(H, A)$ be the fractional space, consisting all $v \in H$ for which the following norm is finite,

$$\|v\|_{H_\alpha} = \|v\|_H + \sup\|\lambda^{1-\alpha} A e^{-\lambda A} v\|_H$$

Theorem 1.6. Assume that $\varphi_N - \sum_{k=1}^p \alpha_k \varphi_{\ell_k} + A\varphi \in H_\alpha$ and (2). Then, problem (10) is well-posed in $C^\alpha(H)$ and the following coercivity estimate holds

$$\begin{aligned} & \left\| \left\{ \tau^{-1} (u_k - u_{k-1}) \right\}_1^N \right\|_{C^\alpha(H)} \\ & + \left\| \left\{ Au_{k-1} \right\}_1^N \right\|_{C^\alpha(H)} + \left\| \left\{ \tau^{-1} (u_k - u_{k-1}) \right\}_1^N \right\|_{C_\tau(H_\alpha)} \\ & \leq M \left(\frac{1}{\alpha} \left\| \phi_N - \sum_{k=1}^p \alpha_k \phi_k + A\phi \right\|_{H_\alpha} + \frac{C(\delta, \theta_p)}{\alpha(1-\alpha)} \|\phi^\tau\|_{C^\alpha(H)} \right), \end{aligned}$$

where M does not depend on φ, φ^τ , and τ .

Proof. Let us establish the estimate for $\left\| \left\{ Au_{k-1} \right\}_1^N \right\|_{C^\alpha(H)}$. Similar arguments introduced in the proof of estimate (24) result that

$$\left\| \left\{ Au_{k-1} \right\}_1^N \right\|_{C_\tau(H)} \leq C(\delta, \theta_p) \left(\frac{1}{\alpha} \|\varphi^\tau\|_{C^\alpha(H)} + \|A\varphi\|_H \right). \tag{36}$$

Next, we estimate

$$\max_{1 \leq k < k+r \leq N} \frac{\|Au_{k-1+r} - Au_{k-1}\|_H}{(r\tau)^\alpha}.$$

Using formula (11), we obtain for $1 \leq k \leq N$ then

$$\begin{aligned}
 Au_{k-1} &= -\varphi_{k-1} + R^{N-k+1}(Au_N + \varphi_N) \\
 &+ \sum_{j=k}^N \tau AR^{j-k+1}(\varphi_j - \varphi_{k-1}) + R^{N-k+1}(\varphi_{k-1} - \varphi_N) \\
 &= J_1(k) + J_2(k) + J_3(k) + J_4(k).
 \end{aligned}
 \tag{37}$$

It is clear that

$$\|J_1\|_{C^\alpha(H)} = \|\varphi^\tau\|_{C^\alpha(H)}.
 \tag{38}$$

Let us estimate $\|J_2\|_{C^\alpha(H)}$. To alleviate the notation, let $v = (Au_N + \varphi_N)$. From the definition of H_α -norm, the equality

$$\sum_{j=k}^{k+r-1} -\tau AR^{N-j} = R^{N-(k-1+r)} - R^{N-k+1}$$

and the formula connecting the resolvent of the generator of a semigroup with the semigroup it follows that

$$\begin{aligned}
 &\|J_2(k+r) - J_2(k)\|_H \\
 &\leq \|v\|_{H_\alpha} \int_0^\infty \sum_{j=k}^{k+r-1} \frac{1}{(N-j-1)!} t^{N-j-1} e^{-t} \frac{e^{-\frac{\pi\delta}{2}t} \tau dt}{\left(\frac{\tau}{2}\right)^{1-\alpha}} \\
 &\leq 2^{1-\alpha} \|v\|_{H_\alpha} (r\tau)^\alpha \frac{2}{\alpha}.
 \end{aligned}
 \tag{39}$$

Thus, using estimate (39), we get

$$\|J_2\|_{C^\alpha(H)} \leq \frac{4}{\alpha} \|v\|_{H_\alpha}.
 \tag{40}$$

It follows from estimate (6), the definition of $C^\alpha(H)$ -norm, and $N\tau = 1$ that

$$\begin{aligned} \|J_3(k)\|_H &\leq \|\varphi^\tau\|_{C^\alpha(H)} \sum_{j=k}^N \frac{\tau}{((j-k+1)\tau)^{1-\alpha}} \\ &\leq \frac{((N-k+1)\tau)^\alpha}{\alpha} \|\varphi^\tau\|_{C^\alpha(H)}, \end{aligned} \tag{41}$$

for all k .

Hence, using estimate (41), we obtain

$$\|J_3(k)\|_H \leq \frac{1}{\alpha} \|\varphi^\tau\|_{C^\alpha(H)}. \tag{42}$$

Next, we estimate

$$\max_{1 \leq k < k+r \leq N} \frac{\|J_3(k+r) - J_3(k)\|_H}{(r\tau)^\alpha}.$$

First, let us consider the case $N - j + 1 \leq 2r$. Using the triangle inequality, estimate (41), we get

$$\frac{\|J_3(k+r) - J_3(k)\|_H}{(r\tau)^\alpha} \leq \frac{(2^\alpha + 1)}{\alpha} \|\varphi^\tau\|_{C^\alpha(H)}. \tag{43}$$

Next, we consider the case $N - j + 1 > 2r$. We can write as

$$J_3(k) - J_3(k+r) = J_{31}(k) + J_{32}(k) + J_{33}(k) + J_{34}(k),$$

where $J_{31}(k) = I_2(t)$, $J_{32}(k) = I_3(k)$, $J_{33}(k) = I_4(k)$ (see equation (26)), and

$$J_{34}(k) = (R^{r-1} - R^{N-(k-1+r)}) (\phi_{k-1+r} - \phi_{k-1}).$$

So we have

$$\|J_{31}(k)\|_H \leq \frac{2^\alpha (r\tau)^\alpha}{\alpha} \|\varphi^\tau\|_{C^\alpha(H)}, \tag{44}$$

$$\|J_{32}(k)\|_H \leq \frac{2^\alpha (r\tau)^\alpha}{\alpha} \|\varphi^\tau\|_{C^\alpha(H)}, \tag{45}$$

$$\|J_{33}(k)\|_H \leq \frac{2^{-1+\alpha} (r\tau)^\alpha}{(1-\alpha)} \|\varphi^\tau\|_{C^\alpha(H)}. \tag{46}$$

Finally, using estimate (5) and the definition of $C^\alpha(H)$ -norm, we get

$$\|J_{34}(k)\|_H \leq 2(r\tau)^\alpha \|\varphi^\tau\|_{C^\alpha(H)}. \tag{47}$$

Hence, it follows from estimates (44)- (47) that for $N - j + 1 > 2r$,

$$\frac{\|J_3(k + \tau) - J_3(k)\|_H}{(r\tau)^\alpha} \leq \frac{M}{\alpha(1-\alpha)} \|\varphi^\tau\|_{C^\alpha(H)}. \tag{48}$$

Combining estimates (43), (48), we get

$$\max_{1 \leq k < k+r \leq N} \frac{\|J_3(k+r) - J_3(k)\|_H}{(r\tau)^\alpha} \leq \frac{M}{\alpha(1-\alpha)} \|\varphi^\tau\|_{C^\alpha(H)}. \tag{49}$$

Thus, estimate (42), (49) result that

$$\|J_3\|_{C^\alpha(H)} = \frac{M}{\alpha(1-\alpha)} \|\varphi^\tau\|_{C^\alpha(H)}. \tag{50}$$

Using estimate (5) and the definition of $C^\alpha(H)$ -norm, we obtain

$$\|J_4(k)\|_H \leq ((N - k + 1)\tau)^\alpha \|\varphi^\tau\|_{C^\alpha(H)} \leq \|\varphi^\tau\|_{C^\alpha(H)}, \text{ for all } k. \tag{51}$$

Hence, estimate (51) gives

$$\|J_4(k)\|_H \leq \|\varphi^\tau\|_{C^\alpha(H)}. \tag{52}$$

By using estimates (5), (7) for $\beta = 0$, we get for all $1 \leq k < k+r \leq N$

$$\begin{aligned} \|J_4(k+r) - J_4(k)\| &\leq \|R^{N-(k-1+r)} - R^{N-k+1}\|_{H \rightarrow H} \|\varphi_{k-1+r} - \varphi_N\|_H \\ &\quad + \|R^{N-(k-1+r)}\|_{H \rightarrow H} \|\varphi_{k-1+r} - \varphi_{k-1}\|_H \\ &\leq (M+1)(r\tau)^\alpha \|\varphi^\tau\|_{C^\alpha(H)}. \end{aligned} \tag{53}$$

So, from estimate (53) it follows that

$$\max_{1 \leq k < k+r \leq N} \frac{\|J_4(k+r) - J_4(k)\|_H}{(r\tau)^\alpha} \leq M_1 \|\varphi^\tau\|_{C^\alpha(H)}. \tag{54}$$

Thus, by combining estimates (52), (54), we obtain

$$\|J_4\|_{C^\alpha(H)} = M_1 \|\varphi^\tau\|_{C^\alpha(H)}. \tag{55}$$

From estimates (36), (38), (40), (50), and (55) it results that

$$\left\| \{Au_{k-1}\}_1^N \right\|_{C^\alpha(H)} \leq M \left(\frac{1}{\alpha} \|Au_N + \varphi_N\|_{H_\alpha} + \frac{1}{\alpha(1-\alpha)} \|\varphi^\tau\|_{C^\alpha(H)} \right). \tag{56}$$

Hence, using the triangle inequality, estimate (56), and difference equation (10), we get

$$\begin{aligned} &\left\| \{\tau^{-1}(u_k - u_{k-1})\}_1^N \right\|_{C^\alpha(H)} \\ &\leq M \left(\frac{1}{\alpha} \|Au_N + \varphi_N\|_{H_\alpha} + \frac{1}{\alpha(1-\alpha)} \|\varphi^\tau\|_{C^\alpha(H)} \right). \end{aligned} \tag{57}$$

Let us now establish the estimate for $\left\| \{\tau^{-1}(u_k - u_{k-1})\}_1^N \right\|_{C^\alpha(H)}$.

It results from formula (11) and difference equation (10) that for all k,

$$\begin{aligned} \frac{u_k - u_{k-1}}{\tau} &= R^{N-(k-1+r)}(Au_N + \varphi_N) + R^{N-(k-1+r)}(\varphi_k - \varphi_N) \\ &\quad - \sum_{j=k}^N \tau AR^{j-(k-1)}(\varphi_j - \varphi_{k-1}) \\ &= G_1(k) + G_2(k) + G_3(k). \end{aligned}$$

Using estimate (5) and the definition of H_α - norm , we obtain

$$\|G_1(k)\|_{H_\alpha} \leq \|Au_N + \varphi_N\|_{H_\alpha}. \tag{58}$$

Now, using the definition of H_α - norm and the formula connecting the resolvent of the generator of a semigroup with the semigroup, we get

$$\begin{aligned} \|G_2(k)\|_{H_\alpha} &= \sup_{\lambda>0} \left\| \lambda^{1-\alpha} A e^{-\lambda A} R^{N-k+1} (\varphi_k - \varphi_N) \right\|_H \\ &= \sup_{\lambda>0} \left\| \lambda^{1-\alpha} A e^{-\lambda A} \int_0^\infty \frac{t^{N-k} e^{-t}}{(N-k)!} e^{-\tau t A} (\varphi_k - \varphi_N) dt \right\|_H \\ &\leq \|\varphi^\tau\|_{C^\alpha(H)}. \end{aligned} \tag{59}$$

Next, let us estimate $\|G_3(k)\|_{H_\alpha}$. Let $\lambda > 0$. From estimates (4), (5), (7) for $\beta = 1$, and identity (20) it follows that

$$\|\tau A e^{-\lambda A} A R^{j-k+1}\|_{H \rightarrow H} \leq \min \left\{ \frac{\tau}{((j-k)\tau)^2}, \frac{\tau}{\lambda^2} \right\} \leq M \frac{\tau}{((j-k)\tau + \lambda)^2}. \tag{60}$$

Using estimate (60) and the definition of $C^\alpha(H)$ - norm , we get

$$\begin{aligned} \|G_3(k)\|_{H_\alpha} &\leq M \|\varphi^\tau\|_{C^\alpha(H)} \sup_{\lambda>0} \lambda^{1-\alpha} \sum_{j=k}^N \frac{\tau}{((j-k)\tau + \lambda)^{2-\alpha}} \\ &\leq M \frac{\|\varphi^\tau\|_{C^\alpha(H)}}{1-\alpha}. \end{aligned} \tag{61}$$

Hence, combining estimates (58)- (61), we obtain

$$\begin{aligned} &\left\| \left\{ \tau^{-1} (u_k - u_{k-1}) \right\}_1^N \right\|_{C^\alpha(H)} \\ &+ \left\| \left\{ Au_k \right\}_1^N \right\|_{C^\alpha(H)} + \left\| \left\{ \tau^{-1} (u_k - u_{k-1}) \right\}_1^N \right\|_{C^\alpha(H)} \\ &\leq M \left(\frac{1}{\alpha} \|Au_N + \varphi_N\|_{H_\alpha} + \frac{1}{\alpha(1-\alpha)} \|\varphi^\tau\|_{C^\alpha(H)} \right). \end{aligned} \tag{63}$$

Now, we estimate $\|Au_N + \varphi_N\|_{H_\alpha}$. Using formula (12), we get

$$\begin{aligned} Au_N + \varphi_N &= T_\tau \left\{ -\sum_{k=1}^p \alpha_k \sum_{j=\ell_{k+1}}^N \tau AR^{j-\ell_k} (\phi_j - \phi_{\ell_k}) \right. \\ &\quad \left. + \sum_{k=1}^p \alpha_k R^{N-\ell_k} (\phi_{\ell_k} - \phi_N) \right. \\ &\quad \left. + \phi_N - \sum_{k=1}^p \alpha_k \phi_{\ell_k} + A\varphi \right\} \\ &= P_1 + P_2 + P_3. \end{aligned}$$

It follows from estimates (6), (9), (60), assumption (2), and the definition of $C^\alpha(H)$ -norm that

$$\|P_1\|_{H_\alpha} \leq \frac{C(\delta, \theta_p)}{(1-\alpha)} \|\varphi^\tau\|_{C^\alpha(H)}, \tag{64}$$

$$\|P_2\|_{H_\alpha} \leq C(\delta, \theta_p) \|\varphi^\tau\|_{C^\alpha(H)}, \tag{65}$$

$$\|P_3\|_{H_\alpha} \leq C(\delta, \theta_p) \left\| \varphi_N - \sum_{k=1}^p \alpha_k \varphi_{\ell_k} + A\varphi \right\|_{H_\alpha}. \tag{66}$$

Therefore, estimates (63), (64)- (66) finishes the proof of Theorem 1.6. □

3 Application

In this section, we consider applications of Theorem 1.5 and Theorem 1.6.

First, let us consider the nonlocal boundary value problem for one dimensional parabolic equation

$$\begin{cases} u_t + (a(x)u_x)_x - \delta_u = f(t,x), & 0 < t < 1, \quad 0 < x < 1, \\ u(1,x) = \sum_{m=1}^p \alpha_m u(\theta_m, x) + \varphi(x), & 0 \leq x \leq 1, \\ 0 \leq \theta_1 < \theta_2 < \dots < \theta_p < 1, \\ u(t,0) = u(t,1), \quad u_x(t,0) = u_x(t,1), & 0 \leq t \leq 1 \end{cases} \quad (67)$$

under assumption (2), where $\delta > 0$, $a(x) \geq a > 0$ ($x \in (0,1)$), $\varphi(x)$ ($x \in [0,1]$)

and $f(t,x)$ ($t,x \in [0,1]$) are smooth functions.

The discretization of problem (67) is carried out in two steps. In the first step, we define the grid space

$$[0,1]_h = \{x = x_n : x_n = nh, \quad 0 \leq n \leq M, \quad Mh = 1\}.$$

Let us introduce the Hilbert space $L_{2h} = L([0,1]_h)$ of the grid functions

$\varphi^h(x) = \{\varphi_n\}_1^{M-1}$ defined on $[0,1]_h$, equipped with the norm

$$\|\varphi^h\|_{L_{2h}} = \left(\sum_{x \in [0,1]_h} |\varphi(x)|^2 h \right)^{\frac{1}{2}}.$$

To the differential operator A generated by problem (67), we assign the difference operator A_h^x by the formula

$$A_h^x \varphi^h(x) = \left\{ - (a(x)\varphi_{\bar{x}})_{x,n} + \delta \varphi_n \right\}_1^{M-1} \quad (68)$$

acting in the space of grid functions $\varphi^h(x) = \{\varphi_n\}_1^{M-1}$ satisfying the conditions

$\varphi_0 = \varphi_M$, $\varphi_1 - \varphi_0 = \varphi_M - \varphi_{M-1}$. It is well-known that A_h^x is a self-adjoint positive definite operator in L_{2h} . With the help of A_h^x , we arrive at the nonlocal boundary value problem

$$\begin{cases} \frac{du^h(t,x)}{dt} - A_h^x u^h(t,x) = f^h(t,x), & 0 < t < 1, \quad x \in [0,1]_h, \\ u^h(1,x) = \sum_{m=1}^p \alpha_m u^h(\theta_m, x) + \varphi(x), & x \in [0,1]_h, \\ 0 \leq \theta_1 < \theta_2 < \dots < \theta_p < 1. \end{cases} \quad (69)$$

In the second step, we replace (69) with the difference scheme (10)

$$\begin{cases} \frac{u_k^h(x) - u_{k-1}^h(x)}{\tau} - A_h^x u_{k-1}^h(x) = f_k^h(x), \\ f_k^h(x) = f^h(t_k, x), \quad t_k = k\tau, \quad 1 \leq k \leq N, \quad x \in [0,1]_h, \\ u_N^h(x) = \sum_{m=1}^p \alpha_m u_{\ell_m}^h(x) + \varphi(x), \quad x \in [0,1]_h, \\ 0 \leq \theta_1 < \theta_2 < \dots < \theta_p < 1. \end{cases} \quad (70)$$

Theorem 2.1. Let τ and h be sufficiently small numbers. Then, the solutions of difference scheme (70) satisfy the following coercivity stability estimate:

$$\begin{aligned} & \left\| \left\{ \tau^{-1} (u_k^h - u_{k-1}^h) \right\}_1^N \right\|_{C_1^\alpha([0,1]_x, L_{2h})} + \left\| \left\{ u_{k-1}^h \right\}_1^N \right\|_{C_1^\alpha([0,1]_x, W_{2h}^2)} \\ & \leq C(\delta, \theta_p) \left(\frac{1}{\alpha(1-\alpha)} \left\| \left\{ f_k^h \right\}_1^N \right\|_{C_1^\alpha([0,1]_x, W_{2h}^2)} + \left\| \varphi^h \right\|_{W_{2h}^2} \right) \end{aligned}$$

hold where $C(\delta, \theta_p)$ is independent of τ , $f_k^h(x)$, and $\varphi^h(x)$, $1 \leq k \leq N - 1$.

Theorem 2.2. Let

$$A_h^x \varphi^h(x) = f_1^h(x) - \sum_{k=1}^p \alpha_k f_{\ell_k}^h(x).$$

Then, for solutions of the problem (70), we have the following stability inequalities

$$\begin{aligned} & \left\| \left\{ \tau^{-1} (u_k^h - u_{k-1}^h) \right\}_1^N \right\|_{C^\alpha([0,1]_r, L_{2h})} + \left\| \left\{ u_{k-1}^h \right\}_1^N \right\|_{C^\alpha([0,1]_r, W_{2h}^2)} \\ & \leq \frac{C(\delta, \theta_p)}{\alpha(1-\alpha)} \left\| \left\{ f_k^h \right\}_1^N \right\|_{C^\alpha(H)}, \end{aligned}$$

where M does not depend on φ, φ^τ , and τ .

The proof of Theorem 2.1, Theorem 2.2 is based on the abstract Theorem 1.5, Theorem 1.6 and the symmetry properties of the difference operator A_h^x defined by formula (68).

Second, let Ω be the unit open cube in the n-dimensional Euclidean space $R^n = \{x = (x_1, \dots, x_n) : 0 < x_i < 1, i = 1, \dots, n\}$ with boundary $S, \bar{\Omega} = \Omega \cup S$. In $[0,1] \times \Omega$, the boundary value problem for the multi-dimensional parabolic equation

$$\begin{cases} \frac{\partial u(t, x)}{\partial t} + \sum_{r=1}^n (a_r(x) u_{x_r})_{x_r} = f(t, x), \\ x = (x_1, \dots, x_n) \in \Omega, \quad 0 < t < 1, \\ u(1, x) = \sum_{i=1}^p \alpha_i u(\theta_i, x) + \varphi(x), \quad x \in \Omega, \\ 0 \leq \theta_1 < \theta_2 < \dots < \theta_p < 1, \\ u(t, x) = 0, \quad x \in S, \quad 0 \leq t \leq 1 \end{cases} \quad (71)$$

under assumption (2) is considered. Here $a_r(x), (x \in \Omega), \varphi(x) (x \in \bar{\Omega})$, and $f(t, x) (t \in (0,1), x \in \Omega)$ are given smooth functions and $a_r(x) \geq a > 0$.

The discretization of problem (71) is carried out in two steps.

In the first step, define the grid space $\tilde{\Omega}_h = \{x = x_m = (h_1 m_1, \dots, h_n m_n) ; m = (m_1, \dots, m_n), 0 \leq m_r \leq N_r, h_r N_r = 1, r = 1, \dots, n\}, \Omega_h = \tilde{\Omega}_h \cap \Omega$,

$$S_h = \tilde{\Omega}_h \cap S.$$

Let L_{2h} denoted the Hilbert space

$$L_{2h} = L_2(\tilde{\Omega}_h) = \left\{ \varphi^h(x) : \left(\sum_{x \in \tilde{\Omega}_h} |\varphi^h(x)|^2 h_1 \cdots h_2 \right)^{\frac{1}{2}} < \infty \right\}.$$

The differential operator A in (71) is replaced with

$$A_h^x u^h(x) = - \sum_{r=1}^n (\alpha_r(x) u_{\bar{x}_r}^h)_{x_r, j_r}, \tag{72}$$

where the difference operator A_h^x is defined on those grid functions $u^h(x) = 0$, for all $x \in S_h$. It is well-known that A_h^x is a self-adjoint positive definite operator in L_{2h} .

Using (71), we get

$$\begin{cases} \frac{du^h(t,x)}{dt} - A_h^x u^h(t,x) = f^h(t,x), & 0 < t < 1, \quad x \in \tilde{\Omega}_h, \\ u^h(1,x) = \sum_{m=1}^p \alpha_m u^h(\theta_m, x) + \varphi(x), & x \in \tilde{\Omega}_h, \\ 0 \leq \theta_1 < \theta_2 < \cdots < \theta_p < 1. \end{cases} \tag{73}$$

From (73) it follows that

$$\begin{cases} \frac{u_k^h(x) - u_{k-1}^h(x)}{\tau} - A_h^x u_{k-1}^h(x) = \varphi_k^h(x), \\ \Omega_k^h(x) = f^h(t_k, x), \quad t_k = k\tau, \quad 1 \leq k \leq N, \quad x \in \tilde{\Omega}_h, \\ u_N^h(x) = \sum_{m=1}^p \alpha_m u_{\theta_m}^h(x) + \varphi(x), \quad x \in \tilde{\Omega}_h, \\ 0 \leq \theta_1 < \theta_2 < \cdots < \theta_p < 1. \end{cases}$$

Theorem 2.3. Let τ and $|h| = \sqrt{h_1^2 + \dots + h_n^2}$ be sufficiently small numbers. Then, the solutions of difference scheme (74) satisfy the following coercivity

$$\begin{aligned} & \left\| \left\{ \tau^{-1} (u_k^h - u_{k-1}^h) \right\}_1^N \right\|_{C_1^\alpha([0,1]_r, L_{2h})} + \left\| \left\{ u_{k-1}^h \right\}_1^N \right\|_{C_1^\alpha([0,1]_r, W_{2h}^2)} \\ & \leq C(\delta, \theta_p) \left(\frac{1}{\alpha(1-\alpha)} \left\| \left\{ f_k^h \right\}_1^N \right\|_{C_1^\alpha([0,1]_r, W_{2h}^2)} + \left\| \phi^h \right\|_{W_{2h}^2} \right), \end{aligned}$$

where $C(\delta, \theta_p)$ is independent of τ , $f_k^h(x)$, and $\varphi^h(x)$, $1 \leq k \leq N - 1$.

Theorem 2.4. Let $A_h^x \varphi^h(x) = \varphi_1^h(x) - \sum_{k=1}^p \alpha_m \varphi_{\ell_m}^h(x)$. Then, for solutions of problem (74), we have the following stability inequalities

$$\begin{aligned} & \left\| \left\{ \tau^{-1} (u_k^h - u_{k-1}^h) \right\}_1^N \right\|_{C^\alpha([0,1]_r, L_{2h})} + \left\| \left\{ u_{k-1}^h \right\}_1^N \right\|_{C^\alpha([0,1]_r, W_{2h}^2)} \\ & \leq \frac{C(\delta, \theta_p)}{\alpha(1-\alpha)} \left\| \left\{ f_k^h \right\}_1^N \right\|_{C^\alpha(H)} \end{aligned}$$

where M does not depend on φ^h , f_k^h , h , and τ .

The proof of Theorem 2.3, Theorem 2.4 is based on the abstract Theorem 1.5, Theorem 1.6, and the symmetry properties of the difference operator A_h^x , defined by formula (72), and the following theorem:

Theorem 2.5. (see Sobolevskii (1975)) For the solutions of the elliptic differential problem

$$\begin{cases} A_h^x u^h(x) = w^h(x), & x \in \tilde{\Omega}_h, \\ u^h(x) = 0, & x \in S_h, \end{cases}$$

the following coercivity inequality holds:

$$\sum_{r=1}^n \left\| (u_k^h)_{\bar{x}_r, \bar{x}_r, j_r} \right\|_{L_{2h}} \leq M \|w^h\|_{L_{2h}}.$$

References

- [1] Aibeche A., Favvini A., Coerciveness estimate for Ventcel boundary value problem for a differential equation , Semigroup Forum, 70, No 2, 269-277, 2005.
- [2] Agarwal R., Bohner M., V.B. , Shakhmurov, Maximal regular boundary value problems in Banach-valued weighted spaces , Bound Value Probl., 1, 9-42, 2005.
- [3] Ashyralyev A., Nonlocal boundary value problems for partial differential equations:well-posedness , AIP Conference Proceedings Global Analysis and Applied Mathematics: International Workshop on Global Analysis, 729, 325-331, 2004.
- [4] Ashyralyev A., Nonlocal boundary value problems for abstract parabolic difference equations: well-posedness in Bochner spaces , J. Evol. Equ., 6, No 1, 1-28, 2006.
- [5] Ashyralyev A., Hanalyev A., Sobolevskii, P.E., Coercive solvability of non local boundary value problem for parabolic equations , Abstr. Appl. Anal., 6, No 1, 53-61, 2001.
- [6] Ashyralyev A., Dural A., Sözen Y., Multipoint nonlocal boundary value problems for reverse parabolic equations: well-posedness , Vestnik of Odessa National University: Mathematics and Mechanics, 13, 1-12, 2008.
- [7] Ashyralyev A., Sobolevskii P.E., Positive operators and the fractional spaces , Offset Laboratory VSU, Voronezh, Russia, 1989.
- [8] Ashyralyev A., Sobolevskii P.E., Well-Posedness of Parabolic Difference Equations, vol. 69 of Operator Theory: Advances and Applications, Birkh"user, Basel, Switzerland, 1994.
- [9] Ashyralyev A., Sobolevskii P.E., New Difference Schemes for Partial Differential Equations, vol. 148 of Operator Theory: Advances and Applications, Birkh"user, Basel, Switzerland, 2004.
- [10] Sobolevskii P.E., Coerciveness inequalities for abstract parabolic equations , Dokl. Akad. Nauk SSSR, 157, No 1, 52-55, 1964.
- [11] Clement Ph., Guerre-Delabriere S., On the regularity of abstract Cauchy

- problems and boundary value problems, *Atti Accad. Naz. Lincei Cl. Sci. Fis. Mat. Natur. Rend. Lincei (9) Mat. Appl.*, 9, No 4, 245-266, 1999.
- [12] Ladyzhenskaya O.A., Solonnikov V.A., Ural'tseva N.N., *Linear and Quasilinear Equations of Parabolic Type*. Moscow: Nauka, 1967, English transl. *Linear and Quasilinear Equations of Parabolic Type*, Providence, RI: American Mathematical Society, *Translations of Mathematical Monographs*, 23, 1968.
- [13] Ladyzhenskaya O.A., Ural'tseva N.N., *Linear and Quasilinear Equations of Elliptic Type*, Moscow: Nauka, 1973 , English transl. of first edition, *Linear and Quasilinear Elliptic Equations*, London, New York: Academic Press, 1968.
- [14] Shakhmurov V.B., Coercive boundary value problems for regular degenerate differential -operator equations, *J. Math. Anal. Appl.*, 292, No 2, 605-620, 2004.
- [15] Sobolevskii P.E., Some properties of the solutions of differential equations in fractional spaces in: *Trudy Nauchn. Issled. Inst. Mat. Voronezh. Gos. Univ.*, 14, 68-74, 1975.
- [16] Sobolevskii P.E., The coercive solvability of difference equations, *Dokl. Acad. Nauk SSSR*, 201-1971, No 5, 1063-1066, 1971.
- [17] Vishik M.L., Myshkis A.D., Oleinik O.A., *Partial Differential Equations in: Mathematics in USSR in the Last 40 Years, 1917-1957*, Moscow: Fizmatgiz, 1,563-599, 1959.